

5 Literaturverzeichnis

- Abusaleh / Dutta, O.P. 1988:** Inter Relationship of yield components in cucumber. Veg. Sci. 15: 79-85.
- Acreman, S.J. / Dixon, A.F.G. 1989:** The effects of temperature and host quality on the rate of increase of the grain aphid (*Sitobion avenae*) on wheat. Annals of applied Biology. 115: 3-9.
- Adams, J. B. / van Emden, H. F. 1972:** The biological properties of Aphids and their host plant relationships. Aphid Technology Ed. van Emden Academic Press, London New York. 48-97.
- Ahmed, M.A. 1994:** Differences in susceptibility of six cucumber cultivars to infestation by *Aphis gossypii* GLOV., *Tetranychus urticae* and *Bemisia tabaci* as correlated to protein and amino acid contents of leaves. Annals of Agricultural Science Moshtohor 32: 2189-2194.
- Albert, R. / Merz, F. 1995:** The cotton aphid is difficult to cope with.- TASPO-Gartenbaumagazin 4: 40-41.
- Aldyhim, Y. N. / Khalil, A. F. 1993:** Influence of temperature and daylength on population development of *Aphis gossypii* on *Cucurbita pepo*. Entomologia experimentalis et Applicata 67: 167-172.
- Amand, P. C. St. / Wehner, T. C. 1991:** Crop loss to 14 diseases in Cucumber in North Carolina for 1983 to 1988. CGC 14-17.
- Anten, N. P. R. / Schieving, F. / Medina, E. / Werger, M. J. / Schuffelen, A. 1995:** Optimal leaf area indices in C3 and C4 mono- and dicotyledonous species at low and high nitrogen availability. Physiologia Plantarum 95, 541-550.
- Asyakin, B.P. 1977:** Effectiveness of the cecidomyiid *Aphidoletes aphidimyza* ROND. (Diptera, Cecidomyiidae) in the control of aphids on vegetable crops in greenhouses. Trudy Vsesoyuznogo Nauchno-Issledovatel'skogo Instituta Zashchity Rastenii, 53: 121-130.
- Attia, A. A. / El-Hamaky, M. A. 1984:** The biology of the cotton aphid *Aphis gossypii* (GLOVER), in Egypt (Homoptera: Aphididae). Bulletin de la Societe Entomologique d'Egypte 65: 359-371.
- Attia, A. A. / El-Hamaky, M. A. 1985:** A survey of the different species of aphids attacking some cucurbit vegetables. Bulletin de la Societe Entomologique d'Egypte 65: 373-381.
- Auclair, J.L. 1959:** Feeding and excretion by the pea aphid, *Acyrtosiphon pisum* (HARR.) (Homoptera: Aphididae), reared on different varieties of peas. Entomologia experimentalis et Applicata. 2: 279-286.
- Auclair, J. L. 1967:** Effects of pH and sucrose on rearing the cotton aphid, *Aphis gossypii*, on a germ-free and holid diet. Journal of Insect Physiology 13: 431-446.
- Auclair, J. L. 1989:** Host Plant Resistance. World Crop Pests. Aphids: their biology, natural enemies and control, Vol. 2C, edited by Minks, A. K. and Harrewijn, P., Elsevier Verlag, 225-254.
- Auclair, J.L., / Maltais, J.B. 1961:** The nitrogen economy of the pea aphid, *Acyrtosiphon pisum* (HARR.), on susceptible and resistant varieties of peas, *Pisum sativum* L. Proceedings of the 11th Int. Congress of Entomology, Vienna, 1960, 1: 740-743.
- Banks, C.J. / Macaulay, E.D. 1965:** The ingestion of nitrogen and solid matter from *Vicia fabae* by *Aphis fabae* SCOP. Annals of applied Biology 55: 207-218.
- Bar-Tal, A. / Feigin, A. / Rylski, I. / Pressman, E. 1994:** Effects of root pruning and N-NO₃ solution concentration on nutrient uptake and transpiration of tomato plants. Scientia Horticulturae 58: 77-90.
- Basavangoud, K. / Kulkarni K.A. / Thontadarya, 1981:** Estimation of crop loss in safflower due to the aphid, *Dactynotus compositae* THEOBLAD (Hemiptera: Aphididae) Mysore Journal of agricultural Science. 15: 279-282.

- Baumeister, W. / Reichart, G. 1969:** Lehrbuch der angewandten Botanik. Gustav Fischer, Stuttgart.
- Bennison, J.A. 1992:** Biological control of aphids on cucumbers use of open rearing systems or "Banker Plants" to aid establishment of *Aphidius matricariae* and *Aphidoletes aphidimyza*. Med. Fac. Landbouww. Univ. Gent, 57: 457-466.
- Beratungsdienst Reichenau e.V. 1998:** Ergebnisse des Sortenversuchs, Sortenspiegel 1998. Unveröffentlicht.
- Bertin, N. / Heuvelink, E. 1993:** Dry-matter in productio in a tomato crop: comparision of two simulation models. Journal of Horticultural Science. 68: 995-1011.
- Birch, L. C. 1948 :** The intrinsic rate of natural increase of an aphid population. Journal of animal Ecology 17: 15-26.
- Blackman, R. L. / Eastop, V. F. 1984:** Aphids on the world's crops. An identification and information guide. John Wiley & Sons, ISBN 0 471904260.
- Blakeney, A. B. / Mutton, L. 1980:** A simple colorimetric method for the determination of sugars in fruit and vegetables. J. Sci. Food Agric. 31: 889-897.
- Bondarenko, N. V. 1987:** The experience of biological and integrated control of pests on glasshouse crops in the USSR. WPRS Bul. X/2: 33-36.
- Bosland, P. W. / Ellington, J. J. 1996:** Comparison of *Capsicum annum* and *C. pubescens* for antixenosis as a means of aphid resistance. Horticultural Science 31: 1017-1018.
- Bouman, B. A. M. 1995:** Crop modelling and remote sensing for yield prediction. Netherlands Journal of Agricultural Science 43: 143-161.
- Bradford, M. M. 1976:** A rapid and sensitive method for quantification of microgram quantities of protein utilizing the priciple of protein-dye binding. Anal. Biochem. 72: 248-254.
- Brinkmann, R. / Oldenburg, S. / Zebitz C. P. W. 1990:** Einfluss saugender Insekten auf Wachstum und Wurzeleistung von Gurke und Ackerbohne. Mitteilungen der BBA 266, 62.
- Brodeur, J. / McNeil, J. 1994:** Seasonal Ecology of *Aphis nigripes* (Hymenoptera: Aphidiidae), a Parasitoid of *Macrosiphum euphorbiae* (Homoptera:Aphididae). Environmental Entomology 23: 292-298.
- Bünger, I. 1996:** Abschlußbericht zum Forschungsprojekt „Umweltschonende Anbauverfahren von Gemüse und Obst“. 15-19. Ministerium für Ländlicher Raum, Landwirtschaft, Ernährung und Forsten (Hrsg.).
- Bünger, I. 1996:** Abschlußbericht zum Forschungsprojekt „Umweltschonende Anbauverfahren von Gemüse und Obst“, Teilgebiet Biologische Schädlingsbekämpfung in Gurkenkulturen unter Glas, 89 Seiten. Ministerium für Ländlicher Raum, Landwirtschaft, Ernährung und Forsten (Hrsg.).
- Bünger, I. 1998:** Abschlußbericht zum Forschungsprojekt „Umweltschonende Anbauverfahren von Gemüse und Obst“. 17-19. Ministerium für Ländlicher Raum, Landwirtschaft, Ernährung und Forsten (Hrsg.).
- Bünger, I. 1999:** Abschlußbericht zum Forschungsprojekt „Umweltschonende Anbauverfahren von Gemüse und Obst“. Teilbereich Biologische Schädlingsbekämpfung in Gewächshausgurken, 56 Seiten. Ministerium für Ländlicher Raum, Landwirtschaft, Ernährung und Forsten (Hrsg.).
- Bünger, I. / Liebig, H.-P. / Zebitz, C.P.W. 1997:** Erfahrungen mit der „Offenen Zucht“ von Blattlaus-Antagonisten bei der Bekämpfung von *Aphis gossypii* GLOVER. Mitt. der Dt. Ges. für allgemeine und angewandte Entomologie 11: 327-330.
- Bünger, I., Liebig, H.-P. / Zebitz, C.P.W. 1999:** Infestation of greenhouse cucumber caused by several species of aphids and their biological control. Gesunde Pflanze 51: 75-80.

- Bünger, I. / Liebig, H.-P. / Zebitz, C.P.W. 2000:** Befalls-Verlust-Relationen zwischen Gewächshausgurken und Blattläusen. 37. Gartenbauwissenschaftliche Tagung, Zürich 107.
- Bünger, I. / Liebig, H.-P. / Zebitz, C.P.W. 2000:** Biologische Bekämpfung verschiedener Blattlausarten an Gewächshausgurke (*Cucumis sativus* L.) in Südwest Deutschland. Tagung zum Pflanzenschutz in Obst und gartenbaulichen Kulturen, Oktober 2000, Temesvar.
- Buysse, J. / Smolder, E. / Merckx, R. 1993:** The role of free sugars and amino acids in the regulation of biomass partitioning and plant growth. *Plant and Soil* 155/156: 191-194.
- Canizares K. A. L. / Goto, R. 1998:** Crescimento e producao de hibridos de pepino em funcao da enxertia. *Horticultura Brasileira* 16: 110-113.
- Capinera, J. L. 1981:** Some effects of infestation by bean aphid, *Aphis fabae* SCOPOLI, on carbohydrate and protein levels in sugarbeet plants, and procedures for estimating economic injury levels. *Zeitschrift für angewandte Entomologie*, 92: 374-384.
- Challa, H. 1990:** Crop growth models for greenhouse climate control. In: Rabbinge, R., Goudriaan, J. Van Keulen, H., Penning de Vries, F. W. T., van Laar H.H. eds. *Theoretical production ecology reflections and perspectives, Simulation Monographs* 34. Wageningen: Pudoc. pp. 125-145.
- Challa, H. 1989:** Modelling for crop growth control. *Acta Horticulturae* 248: 209-216.
- Chamont, S. 1993:** Modelling dry matter allocation in cucumber crops – Competition between fruits and roots. *Acta Horticulturae* 328: 195-203.
- Clarke, N. D. / Shipp, J. L. / Papadopoulos, A. P. / Jarvis, W. R. / Kohsla, S. / Jewett, T. J. / Ferguson, G. 1999 :** Development of the Harrow greenhouse manager: a decision-support system for greenhouse cucumber and tomato. *Computers and Electronics in Agriculture* 24: 195-204.
- Collins, J. K. / Perkins-Veazie, P. 1994:** Resistance in Muskmelon Cultivars to Melon Aphid. *Horticultural Science* 29: 1367.
- Conte, L. 1998:** Neue Perspektive für die Biologische Kontrolle von *Aphis gossypii*. in geschützten Kulturen. *Informatore Fitopatologico* 1-12: 25.
- Conte, L. 1998:** The technique of 'banker plants' for the biological control of *Aphis gossypii* on cucumber. *Informatore Agrario* 54/36: 71-75.
- Conte, L. / Zancanaro, A. / Chiarini, F. A. A. 1999:** Biological Control of *Aphis gossypii* GLOVER in greenhouse cucumbers using 'banker plants' as open rearing units of beneficials: two years of trials. *Culture Protette* 28/7: 63-67.
- Degen, B. 2000:** Gartenbau in Baden-Württemberg. Hrsg.: Ministerium Ländlicher Raum Baden Württemberg.
- Dengler, R. 1991:** Erfahrungen mit der Baumwoll- oder Grünen Gurkenlaus 1990. *Gemüse* 3:153-154.
- deWitt, C. T. 1968:** Model en Theorie. Inaugurele rede, Wageningen Agricultural University, 10. October 1968, 13 Seiten.
- deWitt, C. T. / Rabbinge, R. 1979:** System analysis and dynamic simulation. *EPPO Bull. SROP* 10: 133-142.
- Dhillon, N.P.S. / Wehner T.C. 1991:** Host-plant resistance to insects in cucurbits-germplasm resources, genetics and breeding. *Tropical Pest Management* 37: 421-428.
- Dixon, A. F. G. 1971 a:** The role of aphids in wood formation. *Journal of Applied Ecology* 8: 165-179.
- Dixon, A. F. G. 1976 :** Biologie der Blattläuse. G. Fischer Stuttgart, New York. S. 86.
- Dixon, A. F. G. 1985:** Aphid Ecology. Blacky & Son, New York.

- Dixon, A. F. G. 1987 a:** Parthenogenetic reproduction and the rate of increase in aphids. In: Minks A.K. and Harrewijn P. (eds.): Aphids, pp. 269-287. Elsevier, Amsterdam.
- Dixon, A. F. G. 1987 b:** Evolution and adaptive significance of cyclical parthenogenesis in aphids. In: Minks A.K. and Harrewijn P. (eds.): Aphids, pp. 289-297. Elsevier, Amsterdam.
- Dixon, A. F. G. 1987 c:** Parthenogenetic reproduction and the rate of increase in aphids. Aphids: Their biology, natural enemies and control. World crop pests 2.. Amsterdam, New York: Elsevier. ISBN 0444426302 (B 727.1), pp. 259-287.
- Dixon, A. F. G. / Wratten, S. D. 1971.** Laboratory studies on aggregation size and fecundity in the black bean aphid *Aphis fabae* Scop. Bulletin of Entomological Research 61: 97-111.
- Dufner, J. / Jensen, U. / Schumacher, E. 1992:** Statistik mit SAS. Stuttgart: B.G. Teubner. ISBN 3-519-02088-2.
- Ebert, T. A. / Cartwright, B. 1997:** Biology and Ecology of *A. gossypii* GLOVER (*Homoptera: Aphididae*). Southwestern Entomologist 22: 116-153.
- Edwards, P. J. / Wratten, S. D. 1980:** Ecology of insect-plant interactions. London: Edward Arnold Ltd.
- El-Habi, M. / Sekkat, A. / El Jadd, L. / Boumezzough, A. 2000:** Biology of *Hippodamia variegata* GOEZE (*Col. Coccinellidae*) and its suitability against *A. gossypii* GLOV. (*Hom., Aphididae*) on cucumber under greenhouse conditions. Journal of applied Entomology 124: 365-374.
- El-Habi, M. / Sekkat, A. / El Jadd, L. / Boumezzough, A. 1999:** Dynamique des Populations d'*Aphis gossypii* GLOVER (*Homoptera: Aphididae*) sur Concombre sous serre en zone Côtière Marocaine. Insect Sciences and its Application, 19: 17-22.
- El-Habi, M. / El Jadd, L. / Sekkat, A. / Boumezzough, A. 2000:** Control of *A. gossypii* GLOVER (*Homoptera: Aphididae*) on cucumber under glass by *Coccinella septempunctata* LINNAEUS (*Coleoptera: Coccinellidae*). Insect Sciences and its Application 19: 57-63.
- Van Emden, H.F ed. 1972:** Aphid technology, with special references to the study of Aphids in the field. Academic Press New York.
- Entwistle, J.C. / Dixon, A. F. G. 1987:** Short term forecasting of wheat yield loss caused by the grain aphid (*Sitobion avenae*) in summer. Annals of applied Biology 111: 489-508.
- Förster, H. / Buchenauer, H. / Grossmann, F. 1980:** Nebenwirkungen der systemischen Fungizide Triadimefon und Triadimenol auf Gerstenpflanzen. I Beeinflussung von Wachstum und Ertrag, II Cytokininartige Effekte, III Weitere Beeinflussungen des Stoffwechsels.- Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 87 (8): 473-492, 87 (10): 640-653, 87 (12): 717-730.
- Fouad, S. A. / Abdelmonem, A. E. 1990:** Empirical models for predicting wheat yield in relation to the density levels of the greenbug, *Schizaphis graminum* RONDANI (*Homoptera: Aphididae*). Annals of Agricultural Science, Moshtohor, 28 (3): 1805-1816.
- Freier, B. / Holz, D. / Rossberg, D. / Wenzel, V. / Wetzels, T. 1985:** Darstellung populationsdynamischer Prozesse bei *Macrosiphum avenae* (FABR.) mit Hilfe des Simulationsmodelles PESTSIM-MAC.-MLU, Halle-Wittenberg, Wiss. Beitr. 61: 101-107.
- Freier, B. / Roßberg, D. / Gosselke, U. / Triltsch, H. 2002:** Das Simulationsmodell GETLAUS 01 zur Demonstration von Nützlingseffekten auf Blattläuse. Mitteilungen der DPG 1, 27.
- Freier, B. / Triltsch, H. / Rossberg, D. 1993:** GTLAUS – A model of wheat – cereal aphid – predator interaction and its use in complex agroecological studies. Journal of Plant Diseases and Protection 103: 543-554.
- Freier, B. / Triltsch, H. 1996:** Climate chamber experiments and computer simulations on the influence of increasing temperature on wheat –aphid –predator interactions.- Aspects of Applied Biology 45: 293-298.

- Freier, B. / Trilsch, H. / Rossberg, D. 1996:** GTLAUS- A model of wheat – cereal aphid – predator interaction and its use in complex agroecological studies. Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz 103: 543-554.
- Fritzsche, R. / Dubnik, M. / Podvesler, G. 1974 :** Möglichkeiten der Kombination von Systeminsektiziden zur Vektorbekämpfung mit Mitteln zur Phytophthora- und Kartoffelkäferbekämpfung. Akademie der Landwirtschaftswissenschaften der DDR und Pflanzenschutzamt des Bezirkes Magdeburg.
- Fuchs, W. / Minzenmayer, R. 1995:** Effect of *A. gossypii* in cotton development and yield in west Texas. Southwestern Entomologist. 20: 241-249.
- Furk, C. / Hines, C. M. 1993:** Aspects of insecticide resistance in the melon and cotton aphid *Aphis gossypii*. Plant Pathology.
- Furk, C. / Powell, D. F. / Heyd, S. 1980:** Pirimicarb resistance in the melon and cotton Aphid, *Aphis gossypii* GLOVER. Plant Pathology 29: 191-196.
- Fürst, P. / Pollack, L. / Graser, T. A. / Stehle, P. 1990:** Appraisal of four precolumn derivatization methods for the high performance liquid chromatographic determination of free amino acids in biological materials. Journal of Chromatography, 499: 577-569.
- Gibson, R.W. / Whitehead, D. / Austin, D. J. / Simkins, J. 1976:** Prevention of potato top-roll by aphicide and its effect of leaf area and photosynthesis. Annals of applied Biology 82: 151-153.
- Godel, H. / Graser, T. A. / Földi, P. / Pfänder, P. / Fürst, P. 1984:** Measurement of free amino acids in human biological fluids by high performance liquid chromatography. Journal of Chromatography 297: 49-61
- Gould, H. J. 1971:** Large scale trials of an integrated programme for cucumber pests in commercial nurseries. Plant Pathology 20: 149-156.
- Grafton-Cardwell, E. 1991:** Geographical and Temporal Variation in Response to Insecticides in Various Life Stages of *Aphids gossypii* (Homoptera: Aphididae) Infesting Cotton in California.- Journal of Economic Entomology 84: 741-749.
- Graser, T. A. / Godel, H. G. / Albers, S. / Földi, P. / Fürst, P. 1985:** An ultra rapid and sensitive high performance liquid chromatographic method for determination of tissue and plasma free amino acids. Analytical Biochemistry 151: 142-152.
- Groenendijk, C. A. / van der Werf, W. / Van Dijk E. / Carneiro R. A. 1990:** Modelling the effect as assimilate withdrawel by black bean aphid, *Aphis fabae*, on the growth of sugarbeet plants, infected with beet yellows virus. Med. Fac. Landbouww. Rijksuniv. Gent, 55: 1085-1098.
- Gross, K.C. / Pharr D.M. 1982:** A potential pathway for galactose metabolism in *cucumis sativus* L., a stachyose transporting species. Plant Physiology 69: 117-121.
- Guldmond, J. A. / Tigges, W. T. / de Vrijer, W. F. P. 1994:** Host races of *Aphis gossypii* (Homoptera: Aphididae) on cucumber and chrysanthemum. Environmental Entomology 23: 1235-1240
- Hågvar, B. E. / Hofsvang, T. 1987:** Foraging by the aphid parasitoid *Ephedrus cerasicola* for patchily distributed hosts.- Entomologia experimentalis et Applicata 44: 81-88.
- Hågvar, B. E. / Hofsvang, T. 1991:** Aphid parasitoids (Hymenoptera, Aphididae): biology, host selection and use in biological control. Biocontrol News and Information 12: 13-41.
- Haile, F. J. / Higley, L. G. / Specht, J. E. / Spomer, S. M. 1998:** Soybean leaf morphology and defoliation tolerance. Agron. J. 90: 353-362.
- Hardee, J. 1994:** The response of *Praon spp.* to aphid sex pheromone components in the field. Entomologia experimentalis et Applicata 71: 95-99.

- Heathcote, G. D. 1972:** Evaluating Aphid Population on Plants. In: Van Emden, H. F. (Eds): Aphid technology, with special references to the study of Aphids in the field, 105-145. New York: Academic Press.
- Henneberry, T. J. / Jech, L. F. / de la Torre, T. / Hendrix, D.L. 2000:** Cotton aphid (*Homoptera: Aphididae*) biology, honeydew production, sugar quality and quantity, and relationships to sticky cotton. *Southwestern Entomologist* 25: 161-174.
- Herold, A. 1980:** Regulation of photosynthesis by sink-activity-the missing link. *New Phytol.* 86: 131-144.
- Heuvelink, A. 1996:** Tomato growth and yield: quantitative analysis and synthesis. Thesis, Wageningen – ISBN 90-5485-498-7.
- Hiller, W. 1999:** Abschlußbericht: INTERREG-Programm "Bodensee-Hochrhein", Projekt "Umweltschonende Anbauverfahren von Gemüse und Obst" II, Teilgebiet Gemüsebau. Ed. MLR.
- Holz, F. / Wetzell, T. 1989:** Einschätzung und Nutzung eines Populationsmodells für die Getreideläus *Macrosiphum (Sitobion) avenae* (F.).- *Journal of Applied Entomology.* 108: 328-334.
- Honek, A. / Jaroski, V. / Lapchin, L. / Rabasse, J. M. 1998:** Host choice and offspring allocation in the aphid parasitoid *Aphelinus abdominalis* (Hymenoptera: Aphelinidae). *Journal of Agriculture and Entomology* 15: 209-221.
- Honek, A. / Jaroski, V. / Lapchin, L. / Rabasse, J. M. 1998:** The effect of parasitism by *Aphelinus abdominalis* and drought on the walking movement of aphids. *Entomologia Experimentalis et Applicata* 87: 191-200.
- Horvath, I. / Vigh, L. / van Hasselt, Ph. R. / Woltjes, J. / Kuiper P. J. C. 1983:** Lipid composition in leaves of cucumber genotypes as affected by different temperatures regimes and grafting. *Physiol. Plant.* 57: 532-536.
- Hou, Z. / Chen, Y. Z. / Guo, B. Q. / Yan, F. S. 1997:** EAG and orientation tests on the parasitoid *Lysiphlebia japonica* (Hym., Aphidiidae) to volatile chemicals extracted from host plants of cotton aphid *Aphis gossypii* (Hom., Aphididae). *Journal of Applied Entomology* 121: 495-500.
- Hunt, R. 1982:** Plant growth curves. Edward Arnold. 248 S.
- Hurd, R. G. / Sheard, G. F. 1981:** Fuel saving in greenhouses: the biological aspects. *Grower Guide* 20. London: Grower Books, 55 pp.
- Hurej, M. / van der Werf, W. 1993:** The influence of black bean aphid, *Aphis fabae* SCOP. and its honeydew on leaf growth and dry matter production of sugar beet. *Annals of applied Biology* 122: 201-214.
- Jacobson, R. J. / Croft, P. 1998:** Strategies for the control of *A. gossypii* GLOVER with *A. colemani* HALIDAY in protected cucumbers. *Biocontrol and Science and Technology* 8: 377-387.
- Janoudi, A. K. / Widders, I. E. / Flore, J. A. 1993:** Water deficits and environmental factors affect photosynthesis in leaves of cucumber (*Cucumis sativus*). *Journal of the American Society of Horticultural Sciences* 118: 366-370.
- Johnson, C.G. 1969:** Migration and dispersal of insects by flight. Methuen, London.
- Kandoria, J. L. / Jamwal, R. 1988:** Comparative biology of *Aphis gossypii* on Okra, Brinjal and Chilly in the Punjab, *Indian Journal of Aphidology* 2: 35-39
- Kapatos, E. T. / Stratopoulous, E. T. / Sahinoglou, A. / Tsitsipis, J. A. / Lycouresis, D.P. 1996:** Development of an optimum sampling plan for the population of *A. gossypii* (Hom., Aphididae) on cotton in Greece. *Journal of Applied Entomology* 120: 245-248.
- Van Keulen, H. 1982:** Crop production under semi-arid conditions, as determined by nitrogen and moisture availability. 243-249 in Penning de Vries, F. W. T. and H. H. van Laar, *Simulation Monographs*, Pudoc, Wageningen.

- Khalifa, A. / El Din, S.N. 1964:** Biological and ecological studies on *Aphis gossypii* GLOVER (Homoptera:Aphididae). Bulletin de la Societe Entomologique d'Egypte XLVIII:131-153.
- Kieckhefer, R. W. / Gellner J.L. / Riedell W. 1995:** Integrated Pest Mangement. Evaluation of the aphid-day standard as a predictor of yield loss caused by cereal aphids. Agronomical Journal 87: 785-788.
- Kindlmann, P. / Dixon, A. F. G. 1989:** Developmental constraints in the evolution of reproductive strategies: telescoping of generations in parthenogenetics aphids. Functional Ecology 3: 531-537.
- Kindlmann, P. / Dixon, A. F. G. 1992:** Optimum body size: Effects of food quality and temperature, when reproductive growth rate is restricted, with examples from aphids. Journal of Ecology and Biology 5: 677-690.
- Kindlmann, P. / Dixon, A. F. G. / Gross, L. J. 1992:** The relationship between individual and population growth rates in multicellular organism. Journal of Theoretical Biology 157: 535-542.
- Klingauf, F. A. 1988:** Feeding, Adaption and Excretion. In: Aphids: Their biology, natural enemies and control. World crop pests 2, S. 225-248. Amsterdam, New York: Elsevier. ISBN 0-444-42630-2.
- Klingler, J. / Powell, G. / Thompson G. A. / Isaacs, R. 1998:** Phloem specific aphid resistance in *Cucumis melo* line AR5: effects on feeding behaviour and performance of *Aphis gossypii*. Entomologia Experimentalis et Applicata 86 :79-88.
- Kloft, W. 1960:** Wechselwirkungen zwischen pflanzensaugenden Insekten und den von ihnen besogenen Pflanzengeweben. Habilitationsschrift aus dem Institut für angewandte Zoologie Würzburg. Zeitschrift der angewandten Entomologie. Teil I: 45: 337-381, Teil II: 46: 42-70.
- Kocourek, F. / Beráková, J. / Jarosik, V. 1993:** Introduction of predatory midge *Aphidoletes aphidimyza* (RONDANI) for control of cotton aphid, *Aphis gossypii* GLOVER, on greenhouse cucumbers. Ochrana Rostlin 29/3: 179-185.
- Kocourek, F. / Havelka, J. / Beráková, J. / Jarosik, V. 1994:** Effect of temperature on development rate intrinsic rate of increase of *A. gossypii* reared in greenhouse cucumber. Ochrana Rostlin 71: 59-64.
- Köhler, W. / Schachtel, G. / Voleske, P. 1996:** Biostatistik. Springer Verlag. ISBN 3-540-59252-0.
- Krieg, A. / Franz, J. M. 1989:** Lehrbuch der biologischen Schädlingsbekämpfung. Paul Parey Verlag ISBN 3-489-62326-6.
- Krug, H. 1989:** Growth Models as an Aid for Planning Vegetable Produktion in Greenhouses.- Institute of Vegetable Crops. Acta Horticulturae 260: 391-400.
- Krug, H. 1991:** Gemüseproduktion. Ein Lehr- und Nachschlagwerk für Studium und Praxis. Verlag Paul Parey 2. Auflage; ISBN 3-489-55722-0; 541 p.
- Krug, H. / Liebig, H.-P. 1988:** Static Regression Models for Planning Greenhouse Produktion. Institute for Vegetable Crops. Acta Horticulturae 230: 427-433.
- Krumscheid, B. 1996:** Schaderregerinteraktionen im System Gurke (*Cucumis sativus* L.) – Grüne Gurkenlaus (*Aphis gossypii* GLOVER)- Echter Gurkenmehltau (*Spaerotheca fuliginea* POLL.) Diplomarbeit Universität Hohenheim, 107 Seiten.
- Kunkel, H. / Hertel, R. 1975:** Erstellung einer Stoffwechselbilanz für beide virginoparen Morphen von *Myzus persicae* (SULZ.) – Larven im dritten Stadium. Oecologia (Berlin), 18: 23-35.
- Kuo-Sell, H. L. 1985:** Some bionomics of the predacious aphid midge, *Aphidoletes aphidimyza* (ROND.) (Diptera: Cecidomyiidae), and the possibility of using the rose grain aphid, *Metopolophium dirhodum* (WLK.), as an alternative prey in an open rearing unit in greenhouses. In: Integrated and Biological control in protected crops, pp. 151-161 ed. A.A. Balkema Rotterdam/Netherlands.

- Lapchin, L. / Boll, R. / Rochat, J. / Geria, A.-M. / Franco, E. 1997:** Projection pursuit nonparametric regression used for predicting insect densities from visual abundance classes. *Environmental Entomology* 26: 736-744.
- Leather, S. R / Dixon, A. F. G. 1984:** Aphid growth and reproductive rates. *Entomologia experimentalis et applicata* 35: 137-140.
- Lee, J. H. / Heuvelink, E. / Challa, H. 1998:** Effects of palnting date and plant desity on crop growth of cut chrysanthemum. *Journal of Horticultural Sciences and Biotechnology* 77: 238-247.
- Van Lenteren, J. C. / Woets, J. 1988:** Biological and integrated pest control in greenhouses. *Annual Review of Entomology* 33: 239-269.
- Leski, R. 1995:** Nadkompensacja roslin atakowanych przez szkoniki - Overcompensation of crops injured by pests. In: Pruszyński, S., Lipa, J.J. (Hrsg.): Materials of the 35. Research session of Institute of Plant Protection 1995, 121-131.
- Liebig, H.-P. 1978:** Einflüsse endogener und exogener Faktoren auf die Ertragsbildung von Salatgurken (*Cucumis sativus* L.) unter besonderer Berücksichtigung von Ertragsrhythmik, Bestandesdichte und Schnittmaßnahmen, Dissertation TU Hannover, 155 p.
- Liebig, H.-P. 1982:** Einflüsse endogener und exogener Faktoren auf die Ertragsbildung von Salatgurken (*Cucumis sativus* L.). I. Stoffproduktion und Ertragsrhythmik. *Gartenbauwissenschaft* 1: 4-13.
- Liebig, H.-P. 1983:** Einflüsse endogener und exogener Faktoren auf die Ertragsbildung von Salatgurken. II. Bestandesdichte. *Gartenbauwissenschaft* 2: 81-92.
- Liebig, H.-P. 1989:** Model of Cucumber Growth and Prediction of Yields.- Institute of Vegetable Crops University of Hannover. *Acta Horticulturae* 248: 187-191.
- Liebig, H.-P. 1989:** Development of planning models – methodological aspects. *Acta Horticulturae* 248: 159-165.
- Liebig, H.-P. 1989:** Models to Predict Crop Growth. *Acta Horticulturae* 248: 55-68.
- Liebig, H.-P. 1991:** *Cucurbitaceae* (Kürbisgewächse) in: Gemüseproduktion. Ein Lehr- und Nachschlagwerk für Studium und Praxis. Verlag Paul Parey 2. Auflage; ISBN 3-489-55722-0; 541 p.
- Liebig, H.-P. / Alscher, G. 1993:** Combination of growth models for optimized CO₂- and temperature-control of lettuce. *Acta Horticulturae* 323: 155-162.
- Llwelllyn, M. 1988:** Aphid Energy Budgets. In: Aphids - Their biology, natural enemies and control. *World crop pests* 2, pp. 109-117. Amsterdam, New York: Elsevier. ISBN 0-444-42630-2.
- Lozán, J. L. / Kausch, H. 1998:** Angewandte Statistik für Naturwissenschaftler, 2. Auflage, Pareys Studentexte, 74, ISBN 3-8263-3159-1, Parey Berlin.
- Luczak, I. / Gaweda, M. 1991:** Development of the black bean aphid (*Aphis fabae* Scop.) on red beets in relation to the chemical composition of the leaves. *Folia Horticulturae* III/3: 39-48.
- Luo, Z.-Y. / Shen, X.-Y. 1985:** Preliminary investigation on population dynamics computer simulation and forecast of Cotton Aphid (*Aphis gossypii* GLOVER) in the Sheshan field Shanghai suburb. *Contr. Shanghai Inst. Entomol.*, 5: 67-79.
- MacKenzie 1994:** Agrochemical Service. County National Wes Wood MacKenzie, London
- Madore, M. / Grodzinski, B. 1985:** Effects of CO₂ enrichment on Growth and Phtoassimilate Transport in a Dawrf Cucumber (*Cucumis sativus* L.) Cultivar. *Journal of Plant Physiology* 121: 59-71.
- Mallot, G. A. / Davy, A. J. 1978:** Analysis of the effects of the bird cherry-oat aphid on the growth of barley: unrestricted infestation. *New Phytologist* 80: 209-218.

- Marcelis, L. F. M. 1991:** Effects of sink demand on photosynthesis in cucumber. *Journal of Experimental Botany* 42: 1387-1392.
- Marcelis, L. F. M. 1994:** Fruit growth and dry matter partitioning in cucumber. Thesis Wageningen. ISBN 90-5485-310-7, 173 pp.
- Marcelis, L. F. M. 1994:** Fruit shape in cucumber as influenced by position within the plant, fruit load and temperature. *Scientia Horticulturae* 56: 299-308.
- Marcelis, L. F. M. / Gijzen, H. 1998:** Evaluation under commercial conditions of a model of prediction of the yield of cucumber fruits. *Sc. Hort.* 76: 171-181.
- Marcelis, L. F. M. / Heuvelink, E. / de Koning, A. N. M. 1988:** Dynamic simulation of dry matter distribution in greenhouse crops. *Acta Horticulturae* 248: 269-276.
- Marschner, H. 1993:** Mineral nutrition of higher plants. 2. Auflage. London: Academic press, pp. 889.
- Matitschka, G. 1995:** Vereinfachende Beschreibung der Bruttophotosynthese des physiologisch-dynamischen Simulationsmodells SUCROS und Weiterentwicklung für spezielle Anwendungen im Gemüsebau (Produktionssysteme Kohlrabi und Kopfsalat. Dissertation Ulrich E. Grauer Stuttgart ISBN 3-86186-125-9. 127 S.
- Matsui, M. 1996:** Comparison of the efficacy of three parasites, *Aphidius colemani* Viereck, *Lysiphlebus japonicus* Ashmead and *Aphelinus abdominalis* Dalman, in suppressing the population density of the melon aphid, *A. gossypii* Glover, in plastic cages. *Proceedings of the Kanto-Tosan Plant Protection Society* 43: 203-206.
- Mayr, L. 1973:** Möglichkeiten und Grenzen des Einsatzes von *Aphidoletes aphidimyza* (ROND.) (Diptera, Cecidomyiidae) gegen Blattläuse im Gewächshaus. *Zeitschrift für angewandte Entomologie* 73: 255-260.
- McDonald, A. J. S. / Stadenberg, I. 1993:** Nitrogen supply and the control of expansive growth and functions in leaves and roots. *Plant and Soil* 155/156: 195-198.
- Miles, P. W. 1989:** Specific Responses and Damage caused by Aphidoidea. *World Crop Pests. Aphids: their biology, natural enemies and control*, ed. Minks, A. K. and Harrewijn, P.. Vol. 2C, 23- 43.
- Miles, P. W. 1989:** The Response of Plants to the Feeding of the Aphidoidea: Principles. *World Crop Pests. Aphids: their biology, natural enemies and control*, ed. Minks, A. K. and Harrewijn, P.. Vol. 2C, 1-22.
- Miller, R.A. 1989:** Mechanical stress-induced changes in sugar composition of cell walls from cucumber fruit tissues. *Phytochemistry* 28: 389-392.
- Mineo, G. et al. 1994:** On the common athropods living on winter melon (*Cucurbita melo* L. var. inodorus) in Sicily and economic damage evaluation due to *Aphis gossypii* Glov. (Hom.: Aphididae). *Informatore Fitopatologico* 44: 58-63.
- Minks, A. K. / Harrewijn, P. 1988:** Aphids their biology, natural enemies and control, ed. Elsevier Amsterdam A ISBN 0-444-42630-2, B 0-444-42798-8, C 0-444-42799-6.
- Moll, E. 1996:** Zur Umsetzung biometrischer Verfahren in SAS mit Beispielen aus dem Pflanzenschutz. *Biolog. Bundesanstalt Braunschweig (Hrsg.): Berichte der BBA, Heft 10.* Saphir-Verlag, ISSN 0947-8809.
- Moreno, D. A. / Pulgar, G. / Ruiz, J.M. / Villora, G. / Romero, L. 1998:** Optimum nutrient range in cucumber (*Cucumis sativus* L. cv. Brunex F1 plants): Nitrogen and Phosphorus parameters. *International Journal of Experimental Botany* 63: 191-195., XII.
- Moursi, K. S. / Donia, A. A. / Mesbah, H. A. / Haroun, N. S. 1985:** Comparative biological studies of *Aphis gossypii* Glov. On different host plants. *Annals of Agric. Sc., Moshtohor*, 23: 895-897.
- Muldner, S. / Hoogerbrugge, H. / Altena, K. / Bolckmanns, K. 1999 :** Biological pest control in cucumber in the Netherlands. *IOBC Bull.* 22: 177-180.

- Naveed, M. / Rafique-Attique, M. / Rafique, M. 1995:** Role of leaf hairs of different cotton varieties on the population development of cotton aphid (*Aphis gossypii* GLOVER). Pakistan Journal of Zoology Vol. 27: 277-278.
- Newton, J. D. / Dixon, A. F. G. 1990:** Embryonic growth rate and birth weight of the offspring of apterous and alate aphids: a cost of dispersal. Entomologia experimentalis et applicata 55: 223-229.
- Nordheim, E. V. / Hogg, D. B. / Chen, S.-Y. 1988:** Leslie Matrix Models for Insect Populations with overlapping generations. Estimation and Analysis of Insect Populations ed. McDonald, Manly, Lockwood, Logan Proc. of a Conference held in Laramie, Wyoming, 1988. Springer ISBN 0-387-9698-5, 289-298.
- Nutter, F. N. Jr. 1993:** Terms and Concepts for Yield, Crops Loss, and Disease Thresholds. Plant Disease 211-215.
- O'Brian, P. J. / Graves, J. B. 1992:** Insecticide resistance and reproductive biology of *Aphis gossypii* GLOVER. Southwestern Entomologist 17:115-122.
- Oerke, E.-C. / Steiner, U. 1996:** Ertragsverluste und Pflanzenschutz. Schriftenreihe der Deutschen Phytomedizinischen Gesellschaft, 6, Ulmer Germany, ISBN 3-8001-8917-8
- Ohnesorge, B. 1991:** Tiere als Pflanzenschädlinge. 2. Auflage, Thieme Stuttgart 3-13-535402-04 335 S.
- Owusu, E. O. / Horiike, M. / Hirano, C. 1994:** Population parameters of the cotton aphid *Aphis gossypii*, Glover (Homoptera: Aphididae) infesting cucumber. Journal of Horticultural Science 69: 731-734.
- Owusu, E. O. / Kim, C.-S. / Horiike, M. / Hirano, C. 1996:** Comparative biological and enzymatic studies on host-adapted populations of melon and Cotton Aphid, *Aphis gossypii*. Journal of Agricultural Science 126: 449-453.
- Paddok, F. M. 1919:** The cotton or melon louse: Life history studies. Texas Agricultural Experiment Station, Bulletin 257: 1-54.
- Pennig de Vries, F. W. T. / van Laar, H. H. 1982:** Simulation of growth processes and the model BACROS. In Rabbinge, R., Goudriaan, J. Van Keulen, H., Penning de Vries, F. W. T., van Laar H.H. eds. Theoretical production ecology reflections and perspectives, Simulation Monographs 34. Wageningen: Pudoc. pp. 114-135.
- Pharr, D. M. / Sox, H. N. 1984:** Changes in carbohydrate and enzyme levels during the sink to source transition of leaves of *cucumis sativus* L., A stachyose translocator. Plant Science Letters 35: 184-193.
- Pharr, D. M. / Sox, H. N. / Smart, E. L. / Lower, R. L. 1977 :** Identification and distribution of soluble saccharids in pickling cucumber plants and their fate in fermentation. J. Am. Soc. Hortic. Sci. 102: 406-409.
- Pinna, M. 1992:** Use of *Verticillium lecanii* (Zimm.) for the biological control of *Aphis gossypii* (Glover) on cucumber in protected cultivation. Informatore Fitopatologico; 2:56-58.
- Plaut Z. / Mayoral M.L. / Reinhold L. 1987:** Effect of altered sink : source ratio on photosynthetic metabolism of source leaves. Plant Physiol. 85: 786-791.
- Pöhling, H. M. / Morvan, Y. 1984:** Untersuchungen zur Entwicklung von *Aphis fabae* an resistenten und anfälligen Sorten von *Vicia faba*. Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent 49: 793-802.
- Popov, N. / Zabudskaya, I. A. / Shiiko, E. S. / Mencher E. M. / Shevchenko, V. B. 1992:** Development of glasshouse pests of cucumber under different conditions, in relation to possibilities for their biological control. OEPP EPPO Bulletin 22: 529-536.

- Powell, W. / Wright, A. F. 1988.** The abilities of the parasitoid *Aphidius ervi* HALIDAY and *A. rhopalosiphii* DE STEFANI PEREZ (*Hymenoptera: Braconidae*) to transfer between different known host species and the implications for the use of alternative hosts in pest control strategies. *Bulletin of Entomological Research* 78: 683-693.
- Prüter, C. / Zebitz C. P. W. 1991:** Effects of *Aphis fabae* and *Uromyces viciae-fabae* on the growth of a susceptible and aphid resistant cultivar of *Vicia fabae*. *Annals of applied Biology*, 215-226.
- Rabbinge, R. S. A. / Bastiaans, L. 1989:** Combination models, crop growth and pests and diseases. 217-239. In Rabbinge, R. S. A., Ward, und H. H. van Laar (Eds.): *Simulation Monographs* 32, Pudoc, Wageningen.
- Rabbinge, R. S. A. / deWitt, C. T. 1989:** Systems, models and simulation. 3-15. In Rabbinge, R. S. A., Ward, und H. H. van Laar (Eds.): *Simulation Monographs* 32, Pudoc, Wageningen.
- Rabbinge, R. / Drees, E. M. / van der Graf, M. / Verberne, F.C.M. / Wesselo, A. 1981:** Damage effects of cereal aphids in weath. *Netherlands Journal of Plant Pathology* 87: 217-232.
- Rabbinge, R. / Ward, S. A. / Van Laar, H. H. (eds.) 1989:** *Simulation and system management in crop protection*. Pudoc Wageningen ISBN 90-220-0899-1 (Pp 441.75).
- Ramirez, D. R. / Wehner, T. C. / Miller, C. H. 1988:** Source limitation by defoliation and its effect on dry matter production and yield of cucumber. *Hort Science* 23: 704-706.
- Rascher, B. 1995:** Gurken veredeln mit nematodenresistenten Unterlagen. *Gemüse* 1: 50-51.
- Richardson, P. T. / Baker, D. A. 1984:** Assimilate Transport in Cucurbits. *Journal of Experimental Botany*, 35: 1575-1581.
- Richardson, P. T. / Baker, D. A. / Ho, L. C. 1982:** The chemical composition of cucurbit vascular exsudates. *Journal of experimental Botany*, 33: 1239-1247.
- Richter, O. / Spickermann, U. / Lenz, F. 1991:** A new model of plant growth. *Gartenbauwissenschaften* 56: 99-106.
- Robbins, S. N. / Pharr, D. M. 1987:** Regulation of photosynthetic carbon metabolism in cucumber by light intensity and photosynthetic period. *Plant Physiology* 85: 592-597.
- Robbins, S.N. / Pharr, D. 1988:** Effect of restricted root growth on carbohydrate metabolism and whole plant growth of *Cucumis sativus* L. *Plant Physiology* 87: 409-413.
- Rochat, J. 1997:** Delayed effects in aphid-parasitoid systems: Consequences for evaluating biological control species and their use in augmentation strategies. *Entomophaga* 42: 201-213.
- Rochat, J. / Vanlerberghe-Masutti, F. / Chavigny, P. / Boll, R. / Lapchin, L. 1999:** Inter-strain competition and dispersal in aphids: evidence from a greenhouse study. *Ecological Entomology* 24: 450-464.
- Rossing, W. A. H. / de Groot, J. J. R. / van Roermund, H. J. W. 1989 :** Simulation of aphid damage in winter wheat; a case study. 240-261 In Rabbinge, R. S. A., Ward, und H. H. van Laar (Eds.): *Simulation Monographs* 32, Pudoc, Wageningen.
- Ruiz, J.M. / Romero, L. 1988:** Commercial yield and quality of fruits of cucumber plants cultivated under greenhouse conditions: Response to increases in nitrogen fertilization. *J. Agric. Food Chem.* 46: 4171-4173.
- Rumei, X. / Liying, Z. 1991:** Functional relationship between pest abundance, plant physiology and yield as exemplified by the greenhouse whitefly, *Trialeurodes vaporariorum* Westw. (Hom., Aleyrodidae). *Journal of Applied Entomology* 112: 370-381.
- Ruoching, W. / Weishen, L. 1991:** The source and sink relationship between leaf of different position and fruit during different developing stages in cucumber. *Acta Horticulturae Sinica* 18: 245-250.
- SAS Institute INC. 1988:** *SAS/STAT User's Guide*, Release 6.03 Edition- Cary, NC: SAS Institute Inc.

- Sato, Y., Takada, H. / Katayama, J. 1998:** Evaluation of vinyl house release of *Aphidius colemani* and *Aphidoletes aphidimyza* for controlling cucumber-infesting *A. gossypii*. Scientific Reports of the Kyoto prefectural University Human Environment and Agriculture 50: 75-86.
- Schacht, H. / Schenk, M. 1995:** Steuerung der Düngung von Gewächshausgurken (*Cucumis sativus* L.) in geschlossenen Nährlösungssystemen mit Hilfe eines Simulationsmodells.- Institut für Pflanzenernährung der Universität Hannover.
- Schäller, G. 1961:** Aminosäuren im Speichel und Honigtau der Grünen Apfelblattlaus *Aphis pomi* DEG.-Homoptera. Entomologica Experimentalis et Applicata 4: 73-85.
- Schäller, G. 1963:** Papierchromatographische Analyse der Aminosäuren und Amide des Speichels und Honigtaues von 10 Aphidenarten mit unterschiedlicher Phytopathogenität.- Zool. Jb. Physiol. 70: 399-406.
- Schäller, G. 1968:** Biochemische Analyse des Aphidenspeichels und seine Bedeutung für die Gallenbildung. Zool. Jb. Physiol. Bd. 74: 54-87.
- Schappendonk, A. H. C. M. / Challa, H. / Broekharst, P. W. / Cate, A. J. / Udink, T. 1984:** Dynamic Climate Control; An Optimization Study for Earliness of Cucumber Production.- Scientia Horticulturae 23: 135-150.
- Van Schelt, J. 1994:** The selection and utilisation of parasitoids for aphid control in glasshouses.- Proc. Exper. Appl. Entomol. 5: 217-220.
- Scholze, P. 1992:** Das Körperwachstum bei Blattläusen (Homoptera, Aphididae) als Ausdruck des Nährstoffangebots in der Wirtspflanze. II. Untersuchungen an stielen abgeschnittener Blätter von *Tropaeolum majus* L.- Beitr. Ent. 42.
- Scopes, N. E. A. / Biggerstaff, S. B. 1977:** Natural control of *Aphis gossypii*. Glasshouse crops Research Institute, Annual Report: 96-100.
- Sengonca, C. / Hoffmann, A. / Kleinhenz, B. 1994:** Laboruntersuchungen zur Entwicklung, Lebensdauer und Fruchtbarkeit der Getreideblattlausarten *Sitobion avenae* (F.) und *Rhopalosiphum padi* (L.) (Hom., Aphididae) bei verschiedenen tieferen Temperaturen. Journal of Applied Entomology 117: 224-233.
- Shannag, H. 1989:** Zur Schädigung von Gewächshausgurken, *Cucumis sativus* L., durch die Weiße Fliege, *Trialeurodes vaporariorum* (Westwood). Diss. Uni. Hannover.
- Shannag, H.K. / Thorvilson, H. / El-Shatnawi, M.K. 1998:** Changes in photosynthetic and transpiration rates of cotton leaves infested with the cotton aphid, *Aphis gossypii*: Unrestricted infestation. Annals of applied Biology 132: 13-18.
- Shiiko, E. S. et al. 1991:** *Lysiphlebus* parasite against the melon-field aphid on cucumber. Zashchita Rastenii 12: 42.
- Shipp, J.L. / Wang, K. / Binns, M.R. 1999:** Economic injury levels for western flower thrips on greenhouse cucumber. Integrated Control in Glasshouses IOBC Bulletin. 22: 225-228.
- Slosser, J. E. / Pinchak, W. E. / Frank, W. A. 1992:** Effect of planting date on Cotton Aphid and Bandedwinged Whitefly populations in dryland cotton. Southwestern Entomologist, 17: 89-100.
- Solanki, S. S. / Shah, A. 1989:** Path Analysis of Fruit Yield Components in cucumber Prog. hort. 21: 322-324.
- Srivastava, S. J. 1987:** Nutritional Physiology. Lehrbuch World Crop Pests; Aphids : their biology, natural enemies and control, Volume 2 A, edited by Minks, A. K. and Harrewijn, P., Elsevier Verlag, 99-115.
- Stam, P. A. / Abdelrahman, A. A. / Munir, B. 1994:** Comparison of control action thresholds for *Heliiothis armigera*, *Bemisia tabaci* and *Aphis gossypii* on cotton in the Sudan Gezira and Rahad regions. Crop Protection 13: 503-512.

- Starý, P. 1970:** Biology of Aphid Parasites (*Hymenoptera:Aphidiidae*) with respect to integrated control. Series Entomologica, 6. Dr. W. Junk, N.V.- The Hague: 643 p.
- Starý, P. 1976:** Aphid parasites (*Hymenoptera, Aphidiidae*) of the Mediterranean Area. ed. Dr. W. Junk, B.V., Publishers The Hague ISBN 90 6193 0294.
- Van Steenis, M.J. 1992 a:** Biological control of the cotton aphid, *Aphis gossypii* Glover (Hom., Aphididae) : preintroduction evaluation of natural enemies. Journal of Applied Entomology 114: 362-380.
- Van Steenis, M.J. 1992 b:** Perspectives on biological control of cotton aphid (*Aphis gossypii*) Glover.- Med. Fac. Landbouww. Gent 57: 467-472.
- Van Steenis, M. J. 1993:** Intrinsic rate of increase of *Aphidius colemani* Vier. (Hym., Braconidae), a parasitoid of *Aphis gossypii* Glov. (Hom., Aphididae), at different temperatures. Journal of Applied Entomology 116: 192-198.
- Van Steenis, M.J. 1994.** Evaluation of four Aphidiine Parasitoids (Hym., Braconidae) for Biological Control of *Aphis gossypii* Glov. (Hom., Aphididae). Med. Fac. Landbouww. Univ. Gent. 59/2a.
- Van Steenis, M. J. 1994:** Intrinsic rate of increase of *Lysiphlebus testaceipes* Cresson (Hym., Braconidae) a parasitoid of *Aphis gossypii* Glov. (Hom., Aphididae), at different temperatures. Journal of Applied Entomology 118: 399-406.
- Van Steenis, M. J. / El-Khawass, K. A. M. H. 1994:** Behaviour of *Aphidius colemani* searching for *Aphis gossypii*: Functional response and reaction to previously searched aphid colonies. Biocontrol Science and Technology 5: 339-347.
- Van Steenis, M. J. / El-Khawass, K. A. M. H. / Hemerik, L. / van Lenteren, J. C. 1995:** Time Allocation of the Parasitoid *Aphidius colemani* (Hymenoptera: Aphididae) Foraging for *Aphis gossypii* (Homoptera. Aphidae) on Cucumber Leaves. Journal of Applied Entomology 116: 192-198.
- Van Steenis, M.J. / El-Khawass, K.A.M.H. 1995.** Life history of *Aphis gossypii* on cucumber: influence of temperature, host plant and parasitism. Entomologia experimentalis et Applicata 117: 1-11.
- Storck, H. / Berg, E. 1991:** Produktionsplanung und- führung. in Gemüseproduktion, ed. Krug, H. Paul Parey, 2. Auflage ISBN 3-489-55722-0, 265-275.
- Storer, J. R. / Emden, van H. F. 1995:** Antibiosis and antixenosis of chrysanthemum cultivars to the aphid *Aphis gossypii*. Entomologia Experimentalis et Applicata 77: 307-314.
- Stroosnijder, L. 1982:** Simulation of the soil water balance. 175-192 In Penning de Vries, F. W. T. and H. H. van Laar, Simulation Monographs, Pudoc, Wageningen.
- Técsi, L. I. / Maule, A. J. / Smith, A. M. / Leegood, R. C. 1994 :** Metabolic alterations in cotyledons of *Curcubita pepo* infected by cucumber mosaik virus. Journal of Experimental Botany 45: 1541-1551.
- Titi el, A. 1972:** Die Verteilung der Eier von *Aphidoletes aphidimyza* (ROND.) und ihre Bedeutung für den Einsatz unter Glas. Dis.Univ. Göttingen.
- Titi el, A. 1974:** Auswirkung von der räuberischen Gallmücke *Aphidoletes aphidimyza* (ROND.) (*Itonididae:Diptera*) auf Blattlauspopulationen unter Glas. Zeitschrift für angewandte Entomologie 76: 406-417.
- Tol van, S. / Steenis van, M.J. 1994:** Host preference and host suitability for *Aphidius matricariae* HAL. A. *colmani* VIER. (Hym.: Braconidae), parasitizing *Aphis gossypii* GLOV. and *Myzus persicae* SULZ. (Hom.: Aphididae). Med. Fac. Landb. Univ. Gent 59: 273-280.
- Turner, R. B. 1977:** Quantitative requirements for tyrosine, phenylalanine and tryptophan by the cotton aphid, *Aphis gossypii* GLOVER. Comparative Biochemistry and Physiology 56 A: 203-205.

- Turnipseed, S.G. 1972:** Response of soybeans to foliage losses in south Carolina. *Journal of Economic Entomology* 65: 224-229.
- Ulrichs, C. 2001:** Cotton Aphid, *Aphis gossypii* GLOVER, Sternorrhyncha: Aphididae. <http://www.entomology.de>, ©AVRDC-TUM-CLSU periurban project 2 S.
- Valenzuela, J.L. / Sánchez, A. / Lorente, F. A. / del Río, A. / Romero, L. 1991:** Does N directly influence carbohydrates and pigments in tomato and cucumber? *Acta Horticulturae* 287: 477-483.
- Varley, G.C. / Gradwell, G. R. / Hassell, M.P. 1980:** *Populationsökologie der Insekten*. Thieme Verlag, Stuttgart, N.Y. ISBN 3-13-579501-2
- Veen, W. B. 1985:** Photosynthesis and assimilate transport in potato with top-roll disorder caused by aphid *Macrosiphum euphorbiae*.- *Annals of applied Biology* 107: 319-323.
- Verkleij, F.N. / Challa, H. 1988:** Diurnal export and carbon economy in an expanding source leaf of cucumber at contrasting sources and sink temperature. *Physiologia Plantarum* 74: 284-293.
- Van der Vlugt, J.L.F. 1990:** Plant growth and chemical analysis.- *Norwegian Journal of Agricultural Sciences* 4: 77-90.
- Walch-Liu, P. / Neumann, G. / Bangerth, F. / Engels, C. 2000:** Rapid effects of nitrogen form on leaf morphogenesis in tobacco. *Journal of Experimental Botany* 51: 221-237.
- Ward, S. A. / Rabbinge, R. / Dixon, A. F. G. 1988:** Population development Models. *Aphids Vol 2C*, ed. Minks and Harrewijn, Elsevier 21-30.
- Wardlaw, I. F. 1990:** The control of carbon partitioning in plants. *New Phytol.* 116: 341-381.
- Wareing, P. F. / Patrick, J. 1975:** Source-sink relations and the partitioning of assimilates in the plant. In: *Photosynthesis and productivity in different environments* (J. P. Cooper, ed.) Cambridge Univ. Press, Cambridge, 481-499.
- Watson, D.J. 1952:** The physiological basis of variation in yield. *Adv. Agron.* 4: 101-145.
- Weathersbee, A. A. / Hardee, D. D. 1994:** Abundance of cotton aphids (*Homoptera: Aphididae*) and associated biological control agents on six cotton cultivars. *Journal of Economic Entomology* 87: 258-265.
- Weathersbee, A. A. / Hardee, D. D. / Meredith, W. R. 1995:** Differences in yield response to cotton aphids (*Homoptera: Aphididae*) between smooth-leaf and hairy-leaf isogenic cotton lines. *Journal of Economic Entomology* 88: 749-754.
- Weibull, J. 1988:** Free aminoacids in the phloem sap from oats and barley resistant to *Rhopalosiphum padi*. *Phytochemistry* 27: 2069-2072.
- Weibull, J. / Brishammar, S. / Petterson, J. 1986:** Amino acid analysis of phloem sap from oats and barley: a combination of stylet excision and high performance liquid chromatography. *Entomologia Experimentalis et Applicata* 42: 27-30.
- Weibull, J. / Ronquist, F. / Brishammar, S. 1990:** Free amino acid composition of leaf exudates and phloem sap. *Plant Physiology* 92: 222-226.
- Weidemann, H. L. / Rohloff, H. 1976:** Untersuchungen über Salat- und Gurkenmosaikvirus in Freilandbeständen des Kopfsalates. *Nachrichtenblatt des deutschen Pflanzenschutzdienstes* 28: 106-109.
- Wellings, P.W. / Ward, S. A. / Dixon, A. F. G. / Rabbinge, R. 1989:** Crop loss assessment. *Aphids Vol 2C*, ed. Minks and Harrewijn, Elsevier 49-64.
- Welter, S. C. / Rosenheim, J. A. / Marshall, W. J. / Mau, R. F. L. / Gusukuma-Minuto L. R. 1990:** Effect of *Thrips palmi* and Western Flower Thrips (Thysanoptera: Thripidae) on the Yield, growth, and Carbon Allocation Pattern in cucumbers. *Journal of Economic Entomology* 83: 2092-2101.

- Wennegren, U. / Landin, J. 1992:** Population growth and structure in a variable environment. I. Aphids and temperature variation.- *Oecologia*: 394-405.
- Wien, H. C. 1997:** The Cucumbers: Cucumber, melon, squash and pumpkin. CAB INTERNATIONAL: The Physiology of Vegetable Crops, 345-386.
- Wise, I. L. / Lamb, R. 1995:** Spatial Distribution and Sequential Sampling Methods for the Potato Aphid, *Macrosiphum euphorbiae* (THOMAS) (Homoptera: Aphididae), in Oilseed Flax.- *The Canadian Entomologist* 127: 967-976.
- Wratten, S. D. / Lee, G. / Stevens, D. J. 1979:** Duration of cereal aphid populations and the effect on wheat yield and quality. *Proceedings 1979 BCPC. Pest and Diseases* 1: 1-8.
- Wyatt, I. J. / Brown, S. 1977:** The influence of light intensity, daylength and temperature on increase rates of four glasshouse aphids. *Journal of Applied Ecology*: 391-399.
- Wyatt, I. J. / White, P. F. 1977:** Simple estimation of intrinsic increase rates for aphids and tetranychid mites. *Journal of Applied Ecology*. 14: 757-766.
- Xia, J. Y. / van der Werf, W. / Rabbinge, R. 1999:** Modelling biological control of cotton aphid (*A. gossypii*) by seven spot ladybeetle (*Coccinella septempunctata*) in cotton-wheat relay intercropping systems in northern China. *Environmental Entomology* 28: 307-314.
- Xianyuan, X. / Sterling, W. L. 1987:** Computer simulation of Cotton Aphid population dynamics. *Acta Phytophylacica Sinica* 14: 151-156.
- Yencho, G. et al. 1986:** Economic Injury Level, Action Threshold, and a Yield-loss Modell for the Pea Aphid, *Acyrtosiphon pisum* (Homoptera: Aphididae), on Green Peas, *Pisum sativum*.- *Journal of Economic Entomology* 79: 1681-1687.
- Yu-Chang Liu / Yu-Bing Hwang 1991:** Life table of cotton aphid, *A. gossypii* GLOVER, at various photoperiods. *Chinese Journal of Entomology* 11: 106-116.
- Zebitz, C.P.W. 1991:** Über den modifizierenden Einfluss des Ackerbohnenrostes und anderer Umweltfaktoren auf das Verhältnis zwischen *Vicia faba* (L.) und *Aphis fabae* Scop. Habilitationsschrift Universität Hannover 164 S.
- Zhou, X. / Carter, N. 1989:** A simulation model describing the population dynamics and damage potential of the rose grain aphid, *Metopolophium dirhodum* (WALKER) (Homiptera: Aphididae), in the UK. *Bulletin of Entomological Research* 79: 373-380.
- Zhou, X. / Carter, N. 1992:** Effects of temperature, feeding position and crop growth stage on the population dynamics of the rose grain aphid, *Methopolophium dirhodum* (Homimptera: Aphididae). *Journal of Applied Biology* 27-37.
- Zwölfer, H. / Völkl, W. 1997:** Einfluß des Verhaltens adulter Insekten auf Ressourcen-Nutzung und Populationsdynamik: ein Drei-Komponenten-Modell der Populationsdichte-Steuerung. *Entomologia Generalis* 21(3)

